

Review

Scand J Work Environ Health [2014;40\(1\):5-18](#)

doi:[10.5271/sjweh.3388](#)

The association between long working hours and health: A systematic review of epidemiological evidence

by [Bannai A](#), [Tamakoshi A](#)

This is the first systematic review examining the association between long working hours and health after excluding differences in the definition of long working hours and the influence of shift work. We concluded that long working hours are associated with depression, anxiety, sleep, and coronary heart disease. Further studies dealing with long working hours and shift work appropriately are needed.

Affiliation: Department of Public Health, Graduate School of Medicine, Hokkaido University, Japan, Kita 15 Jo Nishi 7 Chome, Kita-ku, Sapporo-shi, 060-8638, JAPAN. tamaa@med.hokudai.ac.jp

Refers to the following text of the Journal: [2010;36\(2\):85-95](#)

The following article refers to this text: [2015;41\(3\):268-279](#)

Key terms: [all-cause mortality](#); [anxiety](#); [circulatory disease](#); [depression](#); [diabetes mellitus](#); [epidemiological evidence](#); [health](#); [health-related behavior](#); [long working hour](#); [overtime](#); [review](#); [shift work](#); [sleep](#); [systematic review](#); [working hour](#); [working time](#)

Additional material

Please note that there is additional material available belonging to this article on the [Scandinavian Journal of Work, Environment & Health -website](#).

The association between long working hours and health: A systematic review of epidemiological evidence

by Akira Bannai, MD,¹ Akiko Tamakoshi, MD, PhD¹

Bannai A, Tamakoshi A. The association between long working hours and health: A systematic review of epidemiological evidence. *Scand J Work Environ Health*. 2014;40(1):5–18. doi:10.5271/sjweh.3388

Objectives Many studies have investigated the association between long working hours and health. By focusing on differences in the definition of long working hours and the influence of shift work, we attempt to explain why the results of these studies remain inconclusive.

Methods We defined long working hours as working time greater than around 40 hours per week or 8 hours per day. Since previous studies have indicated that shift work is detrimental to health, we minimized the influence of shift work in the studies. We also placed importance on the existence of reference groups since this made the results clearer. Based on these points, we analyzed previous studies to clarify the epidemiological evidence regarding the association between long working hours and health. We established inclusion criteria and carried out a systematic search for articles published in the Medline and PsycINFO databases between 1995–2012.

Results We identified a total of 17 articles and 19 studies (12 prospective cohort and 7 cross-sectional studies). The outcomes were all-cause mortality, circulatory disease, diabetes mellitus, metabolic syndrome, depressive state, anxiety, other psychological disorders, sleep condition, cognitive function, and health-related behavior. Long working hours had significant adverse effects on most health outcomes.

Conclusions We concluded that working long hours is associated with depressive state, anxiety, sleep condition, and coronary heart disease. However, further studies that appropriately deal with the definition of long working hours and shift work are needed.

Key terms all-cause mortality; anxiety; circulatory disease; depression; diabetes mellitus; health-related behavior; overtime; shift work; sleep; working time.

In Japan, long working hours is an issue that needs to be addressed promptly for the sake of workers' health. *Karoshi* (sudden death caused by cardiovascular or cerebrovascular disease due to overwork) and *karojisatsu* (suicide due to overwork) may also be related to working long hours. According to data from the Japanese Ministry of Health, Labor, and Welfare, the number of workers suffering from cardiovascular disease, cerebrovascular disease, and mental disorders due to work has increased by about threefold in the last decade (1, 2). Long working hours are common in many countries. According to statistics from the Organisation for Economic Cooperation and Development (OECD) on annual average working hours (3), Korea had the longest working hours between 1980–2007. However, in 2008, Korea

dropped to second place behind Mexico. Similar to Japan, *karoshi* became recognized as a problem in Korea from the early 1990s (4) and consequently the number of working hours may deliberately have been reduced. An International Labor Organization (ILO) report (5) found that the proportion of workers working ≥ 49 or 50 hours/week in 2004–2005 was 49.5% in Korea, 23.6% in New Zealand, 20.4% in Australia, 18.1% in the US, and 14.7% in France. It also estimated that 22.0% of workers globally were working >48 hours/week.

Based on such data, many studies have examined the association between long working hours and health. Various outcomes such as all-cause mortality (6), disease (especially cardiovascular disease) (6, 7), biological indices [heart rate variability, blood pressure (BP),

¹ Department of Public Health, Graduate School of Medicine, Hokkaido University, Sapporo, Japan.

Correspondence to: Akiko Tamakoshi, Department of Public Health, Graduate School of Medicine, Hokkaido University, Japan, Kita 15 Jo Nishi 7 Chome, Kita-ku, Sapporo-shi, 060-8638, JAPAN. [E-mail: tamaa@med.hokudai.ac.jp]

respiratory sinus arrhythmia, etc] (8–10), sleep (8, 9), depressive state (11), alcohol use (12), body mass index (8–10), fatigue (8–10), and general health status (13) have been investigated and positive, negative, or no association reported. The influence of long working hours on human health remains controversial, and we suggest that inconsistencies in the results are attributable to several factors such as the definition of long working hours, characteristics of participants, inclusion of shift work, measurement of outcomes, and potential covariates. In this systematic review, we mainly focused on excluding differences in the definition of long working hours and the influence of shift work.

Long working hours is recognized as working for a length of time which exceeds standard working hours. However, the definition of standard working hours may differ from country to country. For example, standard working hours are 35, 37, and 40 hours/week in France (14), Denmark (15) and the US (14), respectively. In Japan, the Labor Standards Law defines the maximum working time as 40 hours/week. The ILO Convention “Hours of Work to 40 a Week (C047)” was adopted in 1935 (16), but most countries, including Japan, have not ratified it. Under the European Union’s Working Time Directive (2003/88/EC), worktime, including overtime, was limited to 48 hours/week. The difference in standard working hours might affect the definition of long working hours in studies examining the association between long working hours and health. Consequently, this might influence the results of some studies. Therefore, we recognize that standard working hours are around 40 hours/week or 8 hours/day, and long working hours are defined as working longer than this. Moreover, in a study which investigates the association between long working hours and health, we believe that the results would be clearer and easier to understand when standard working time is used as the reference category in the analysis.

There are many types of shift work schedules, for example, night, irregular, and rotating shifts. Many studies have elucidated the association between shift work and health (17–22), and shift work is considered detrimental to health. Shift work has been associated with increased risk for myocardial infarction, coronary events, ischemic stroke (17), gastrointestinal symptoms, gastric ulcers (18), and metabolic syndrome (19). Female night-shift workers have also been reported to be at increased risk of breast cancer (20). Therefore, we must be careful when interpreting the results of studies with shift workers that investigate the association between working long hours and health. We believe that the statistical analyses in studies with shift workers need to be adjusted, by either excluding the shift workers from the analysis, stratifying the data according to work schedules, or adjusting for work schedules as covariates.

The purpose of this paper was to clarify the epide-

miological evidence regarding the association between long working hours and health from previous studies by excluding differences in the definition of long working hours and the detrimental effect of shift work. Accordingly, we set our inclusion criteria and reviewed previous studies.

Methods

Search procedure

We conducted a systematic search of articles published in Medline and PsycINFO between 1995–2012 using the search terms: “work hours”, “workhours”, “working hours”, “long hours”, “extended hours”, “overtime”, “overwork”, and “extended work”. Articles had to be published in English and peer-reviewed, with an abstract available. In the initial search, abstracts were checked for the inclusion criteria (see below). Next, fulltexts of the included articles were obtained and a detailed evaluation performed. The principal author conducted both the initial search and a detailed evaluation of the fulltext twice with an interval in between evaluations. Both authors analyzed those articles that finally remained.

Inclusion criteria

Study type. We included observational studies, such as prospective cohort, case–control, and cross-sectional studies. We excluded experimental and case studies, systematic reviews, meta-analyses, letters, and comments.

Participants. We included articles that did not incorporate shift workers, unless the statistical analysis was performed either by excluding shift workers, stratifying according to work schedules, or adjusting for work schedules as covariates. As described previously, we required a reference group that included participants working around 40 hours/week (8 hours/day) (ie, standard working time).

Working hours. The definition of working hours was time spent on work. Thus, we included hours of work brought home into the total working hours. The reason for this is that participants who worked outside the home but engaged in housework might go home with their work at a regular time (eg, 17:00 hours) but continue working at home. Working hours had to be described as working hours per day, week, or month. Otherwise, articles needed to indicate normal working hours (eg, working hours of contract) and the number of hours overtime. Similar to a previous review (23), we selected articles if the average working hours of participants

exceeded 40 hours/week (8 hours/day), or average working hours plus one standard deviation (SD) were ≥ 40 hours/week (8 hours/day). This was because if the reference group included many participants with extremely short working hours, the effect of long working hours on health might be overestimated. Articles that covered compressed work weeks and temporarily busy periods were excluded.

Outcomes. The World Health Organization's (WHO's) definition of "health" is broad. In this review, we focused on outcomes that directly cause death or disease (eg, diagnosed diseases, sleep problems, depressive state or other psychologically disorders, and health-damaging behavior such as smoking). We excluded job satisfaction, job involvement, work-family conflict, well-being, quality of life, and work performance. We did not cover outcomes related to stress, such as the effort-reward imbalance or demand-control models and burnout.

Accidents, injuries, and specific symptoms or illnesses related to work procedures were also excluded, because such outcomes might be specific to the job (eg, shoulder stiffness as a result of using visual display terminals).

Results

We identified a total of 5088 articles published in English and peer-reviewed with abstracts (figure 1). After excluding duplicates and reviewing abstracts, a detailed evaluation of the fulltext of 149 articles was performed. Finally, after taking inclusion criteria into consideration, 17 articles remained. Both authors examined these articles for details such as study design, characteristics of participants, working hours, definition and measurement of outcomes, statistical analysis, covariates, and results. Any disagreements were solved through discus-




Figure 1. Flow diagram of search procedures.

Table 1. Category and number of outcomes from the final 17 remaining articles based on inclusion criteria. [BP=blood pressure; IFG=impaired fasting glucose]

Category	Outcomes
All-cause mortality	1
Circulatory disease (including change of BP)	6
Diabetes mellitus (including IFG)	4
Metabolic syndrome	1
Mental state	9
Depressive state	5
Anxiety	2
Others	2
Sleep	5
Cognitive function	1
Behavior	5
Alcohol	2
Physical activity	1
Smoking	1
Weight gain	1

sion. When necessary, the authors of the articles were contacted.

In total, 19 studies (12 prospective cohort and 7 cross-sectional studies) were identified from the 17 articles, and a total of 32 outcomes are presented in table 1. The 19 studies are summarized briefly in tables 2 and 3. They are described in more detail in online Appendices A and B (http://www.sjweh.fi/data_repository.php).

Prospective cohort design

Nakamura et al (24) investigated the association between long working hours and one-year change in systolic and diastolic blood pressure (BP) among workers from the manufacturing industry. The average monthly hours of overtime were calculated from recorded timecards. Significant results were observed only among assembly-line workers. The one-year change in diastolic BP in the group working ≥ 80.0 hours/month of overtime was 5.3 mm Hg [95% confidence interval (95% CI) 2.7–7.9], which was significantly increased compared to that of the reference group [1.5 mm Hg (95% CI 0.8–2.2)].

Virtanen et al reported five articles (and six studies) (25–29) based on data from the Whitehall II study in the UK, which was established in 1985 with participants recruited from 20 London-based Civil Service departments. For convenience, we labeled the articles: Study A, 2012 (25), Study B, 2011 (26), Study C, 2010 (27), Studies D and E, June 2009 (28), and Study F, March 2009 (29). Since the article published in June 2009 (28) reported one cohort study with two different measurements for working hours, we treated this as two different studies (studies D and E). The measurement point of working hours was only at baseline in study D but twice in study E.

Study A (25) investigated the association between long working hours and the onset of major depressive episode (MDE) in the preceding 12-month period. Participants were followed for an average of 5.8 [standard deviation (SD) 0.4] years. MDE was evaluated using the University of Michigan's version of the Composite International Diagnostic Interview (UM-CIDI), and the incidence was 3.1%. Participants in the group working 11–12 hours/day had a significantly increased risk of MDE [adjusted odds ratio (OR_{adj}) 2.52 (95% CI 1.12–5.65)] compared to the group working 7–8 hours/day.

Study B (26) evaluated the effect of long working hours on mental health (depressive and anxiety symptoms). The mean follow-up period was 5.3 (SD 0.9) years for depressive symptoms and 5.2 (SD 0.9) years for anxiety symptoms. Outcomes were evaluated by subscales of the General Health Questionnaire (GHQ). The incidence was 2.0/100 person-years for depressive symptoms and 2.3/100 person-years for anxiety symptoms. The adjusted hazard ratio (HR_{adj}) of participants in the group working >55 hours/week was significantly increased at 1.66 (95% CI 1.06–2.61) for depressive symptoms and 1.74 (95% CI 1.15–2.61) for anxiety symptoms compared to the group working 35–40 hours/week. Stratified analysis by gender was also conducted because of a significant interaction between working hours and gender. The significant results were one-sided for women.

Study C (27) was performed to investigate the association between long working hours and incidence of coronary heart disease (CHD) and all-cause mortality. The incidence of CHD included coronary death, incidence of non-fatal myocardial infarction, or incidence of definite angina pectoris. Participants were followed for an average of 11.2 (SD 2.7) years. The incidence of CHD was 5.46/1000 person-years. Participants in the group working 11–12 hours/day had a significantly increased risk [HR_{adj} 1.56 (95% CI 1.11–2.19)] of developing CHD compared to the reference group. A similar increased risk was found for the outcome limited to coronary death or incidence of non-fatal myocardial infarction in the same group [HR_{adj} 1.67 (95% CI 1.02–2.76)]. Long working hours were not significantly associated with all-cause mortality.

Studies D (28) and E (28) evaluated the association between long working hours and sleep condition. In study E, working hours were measured twice in 1991–1994 (before baseline) and 1997–1999 (baseline), and mean working hours were used in the statistical analysis. Outcomes were short sleeping hours, difficulty in falling asleep, frequent waking during the night, early waking, and waking without feeling refreshed. Participants in the group working >55 hours/week were at significantly increased risk for having difficulty in falling asleep: OR_{adj} 4.12 (95% CI 1.71–9.94) and OR_{adj} 7.94 (95% CI 2.97–21.25) in studies D and E, respectively. There were more significant associations in study E compared to D.

Table 2. Characteristics of 12 prospective cohort studies. [ADA=American Diabetes Association; ANCOVA=analysis of covariance; ANOVA=analysis of variance; CHD=coronary heart disease; CIDI=Composite International Diagnostic Interview; CoxPH=Cox proportional hazard analysis; dBP=diastolic blood pressure; DM=diabetes mellitus; GHQ=General Health Questionnaire; h/d=hours per day; h/m=hours per month; h/w=hours per week; ICD-10=International Statistical Classification of Diseases and Related Health Behavior, tenth revision; IFG=impaired fasting glucose; LogRA=logistic regression analysis; MANCOVA=multiple analysis of covariance; MDE=major depressive episode; MI=myocardial infarction; ns=no association; sBP=systolic blood pressure; SD=standard deviation; Sig=significant; UM-CIDI=University of Michigan's version of the Composite International Diagnostic Interview; WHO=World Health Organization; (+)=positive association; (-)=negative association]

Author	Participants	Baseline → Follow-up	Outcomes (measures), Statistical analysis	Working hours	Results
Nakamura et al, 2012 (24)	N=1235, Japan, 611 assembly-line workers, 315 clerks and 309 engineers/special technicians in light metal products manufacturing factory, all male, age 39.3 (SD=10.2, range 20–59) years old	2004 → 2005	1-year change of sBP and dBP (automatic manometer), ANCOVA, multiple post-hoc comparison (Bonferroni correction)	Number of overtime <40.0 ^a , 40.0–79.9, ≥80.0 h/m (scheduled working hours; 40 h/w)	Assembly-line workers: sBP: ns, dBP: Sig, 1.5 mm Hg (reference), 2.3 mm Hg (overtime 40.0–79.9 h/m), 5.3 mm Hg (overtime ≥80.0 h/m, Sig (+) compared with reference) Clerks: sBP and dBP: ns Engineers/special technicians: sBP and dBP: ns
Virtanen et al, 2012 (25): study A	N=2123, UK, civil servants, male 77%, mean age 46.7 years old	1991–1993 (phase 3) → 1997–1999 (phase 5)	Major depressive episode (UM-CIDI), LogRA	Working 7–8 ^a , 9, 10, 11–12 h/d	Sig (+): 2.52-fold (working 11–12 h/d)
Virtanen et al, 2011 (26): study B	Basic participants: N=2960, UK, civil servants, male 76%, mean age 55.3 years old, eligible for depressive symptoms: N=2549, anxiety symptoms: N=2618	1997–1999 (phase 5) → 2001 (phase 6) and 2002–2004 (phase 7)	Depressive symptoms, anxiety symptoms (questions derived from GHQ), CoxPH	Working 35–40 ^a , 41–55, >55 h/w	Total analysis Depressive symptoms: Sig (+): 1.66-fold (working >55 h/w) (linear trend ^b Sig (+): 1.17-fold) Anxiety symptoms: Sig (+): 1.74-fold (working >55 h/w) (linear trend ^b Sig (+): 1.22-fold) Men only Depressive symptoms: ns (linear trend ^b ns) Anxiety symptoms: ns (linear trend ^b Sig (+): 1.19-fold) Women only Depressive symptoms: Sig (+): 2.15-fold (working 41–55 h/w) 2.67-fold (working >55 h/w) (linear trend ^b Sig (+): 1.40-fold) Anxiety symptoms: Sig (+): 1.69-fold (working 41–55 h/w) 2.84-fold (working >55 h/w) (linear trend ^b Sig (+): 1.31-fold)
Virtanen et al, 2010 (27): study C	N=6014, UK, civil servants, male 71%, mean age 48.7 years old (range 39–61)	1991–1994 (phase 3) → 2002–2004 (phase 7)	CHD, coronary death or incidence of non-fatal MI, all-cause mortality (death certificate, electrocardiograms and cardiac enzymes etc, clinical records and use of nitrate medication), CoxPH	Working 7–8 ^a , 9, 10, 11–12 h/d	CHD: Sig (+): 1.56-fold (working 11–12 h/d) Incidence of coronary death or non-fatal myocardial infarction: Sig(+): 1.67-fold (working 11–12 h/d) All-cause mortality: ns
Virtanen et al, June 2009 (28): study D	N=937–1594, UK, civil servants, male 76%, mean age 52.2 years old	1997–1999 (phase 5) → 2002–2004 (phase 7)	Sleep (short sleeping hours (<7 hours/day), difficulty in falling asleep, frequent waking during the night, early waking, waking without feeling refreshed) (self-reported, Jenkins scale), LogRA	Working 35–40 ^a , 41–55, >55 h/w	Short sleeping hours: ns (linear trend ^b Sig (+): 1.30-fold) Difficulty falling asleep: Sig (+): 4.12-fold (working >55 h/w) (linear trend ^b Sig(+): 1.57-fold) Frequent waking: ns (linear trend ^b ns) Early waking: ns (linear trend ^b ns) Waking unrefreshed: ns (linear trend ^b ns)
Virtanen et al, June 2009 (28): study E	N=886–1510, U.K., civil servants, male 76%, mean age 52.2 years old	1997–1999 (phase 5) → 2002–2004 (phase 7)	Sleep (short sleeping hours (<7 hours/day), difficulty in falling asleep, frequent waking during the night, early waking, waking without feeling refreshed) (self-reported, Jenkins scale), LogRA (working hours were measured in 1991–1994 and 1997–1999)	Working 35–40 ^a , 41–55, >55 h/w	Short sleeping hours: Sig (+): 1.55-fold (working 41–55 h/w), 2.80-fold (working >55 h/w) (linear trend ^b Sig (+): 1.56-fold) Difficulty in falling asleep: Sig (+): 7.94-fold (working >55 h/w) (linear trend ^b Sig(+): 2.06-fold) Frequent waking: ns (linear trend ^b ns) Early waking: Sig (+): 2.03-fold (working >55 h/w) (linear trend ^b ns) Waking unrefreshed: ns (linear trend ^b ns)

Continued

Table 2. Continued

Author	Participants	Baseline → Follow-up	Outcomes (measures), Statistical analysis	Working hours	Results
Virtanen et al, March 2009 (29): study F	N=2214, UK, civil servants, male 77%, mean age 52.1 (SD=4.2, range 45–66) years old	1997–1999 (phase 5)→2002–2004 (phase 7)	Cognitive function: high scores represent good cognitive function (short-term memory test, Alice Heim 4-I test (reasoning test), Mill Hill vocabulary test, phonemic fluency test, semantic fluency test), MANCOVA, ANOVA	Working 35–40 ^a , 41–55, >55 h/w	Scores at baseline Mill Hill vocabulary test: Sig (-): working 41–55 h/w, >55 h/w (linear trend ^c Sig (-)) Phonemic fluency test: ns (linear trend ^c Sig(+)) Other three tests: ns Scores at follow-up Alice Heim 4-I test score: Sig (-): working >55 h/w (linear trend ^c Sig (-)) Mill Hill vocabulary test: Sig (-): working 41–55 h/w, >55 h/w (linear trend ^c Sig (-)) Other three tests: ns Change in Alice Heim 4-I test score between baseline and follow-up: Sig (-): reference (-1.77), working 41–55 h/w (-2.23), working >55 h/w (-2.90) (linear trend ^c Sig (-))
Tarumi et al, 2003 (30)	N=867 (mental disorders) and 824 (circulatory disorders), Japan, white-collar office workers in manufacturing company, age 20–60 years old	1997–2000→2001	Mental disorders (ICD-10: F00–F99, G90), circulatory disorders (ICD-10: I00–99) (medical insurance claim records), CoxPH	Working <45 ^a , ≥45 h/w	Mental disorders: ns Circulatory disorders: ns
Nakanishi et al, September 2001 (31)	N=1266, Japan, office workers, all male, age 35–59 years old	1994→1999	IFG or type 2 DM, IFG, type 2 DM (ADA criteria (1997): IFG: fasting plasma glucose concentration of 6.1–6.9 mmol/l, type 2 DM: fasting plasma glucose concentration of ≥7.0 mmol/l or taking hypoglycemic medication), CoxPH	Working <8.0 ^a , 8.0–8.9, 9.0–9.9, 10.0–10.9, ≥11.0 h/d	IFG or type 2 DM: Sig (-): 0.50-fold (working ≥11.0 h/d) (linear trend ^d Sig (-)) IFG: ns Type 2 DM: Sig (-): 0.30-fold (working ≥11.0 h/d) (linear trend ^d Sig (-))
Nakanishi et al, May 2001 (32)	N=941, Japan, white-collar workers, all male, age 35–54 years old	1994→1999	Hypertension (WHO criteria (1978): sBP ≥160 mm Hg and/or DBP ≥95 mm Hg or taking antihypertensive medications), borderline hypertension (sphygmomanometer), CoxPH	Working <8.0 ^a , 8.0–8.9, 9.0–9.9, 10.0–10.9, ≥11.0 h/d	Hypertension: Sig (-): 0.33-fold (working ≥11.0 h/d) (linear trend ^d Sig (-)) Borderline hypertension: Sig (-) 0.63-fold (working 10.0–10.9 h/d), 0.48-fold (working ≥11.0 h/d) (linear trend ^d Sig (-))
Kawakami et al, 1999 (33)	N=2194, Japan, industrial workers in electrical company, all male, including shift workers (46.3%)	1984→1992	Non-insulin dependent diabetes mellitus (NIDDM) (WHO criteria, 1980) – see the details in the text, CoxPH	Number of over-time 0–25 ^a , 26–50, >50 h/m (working hours of contract; 40 h/w)	NIDDM (type 2 DM): Sig (+) 3.73-fold (overtime >50 h/m) (linear trend ^d Sig (+))
Shields, 1999 (34)	N=3830, Canada, various job, male 57%, age 25–54 years old, including shift workers (19.7%)	1994/1995 →1996/1997	Major depressive episode (CIDI), unhealthy weight gain, increased daily smoking, increased alcohol consumption, decreased physical activity, LogRA	Working ≥35 and <41 h/w (standard), ≥41 h/w (long) ^a standard at baseline for MDE, standard-standard pattern for other outcomes	Major depressive episode Men: ns Women: Sig (+) 2.2-fold (working ≥41 h/w) Unhealthy weight gain Men: ns (long-long) Women: ns (long-long) Increased daily smoking Men: ns (long-long) Women: ns (long-long) Increased alcohol consumption Men: ns (long-long) Women: ns (long-long) Decreased physical activity Men: ns (long-long) Women: ns (long-long)

^a Reference working group.^b Linear trend=increasing 10-hour working time (per 10-hour increase).^c Linear trend=entering total working hours into the model as continuous variable.^d Linear trend=increasing categories of working hours per day or month.

Table 3. Characteristics of seven cross-sectional studies. [AUDIT=Alcohol Use Disorders Identification Test; CFSI=Cumulative Fatigue Symptom Index; GHQ=General Health Questionnaire; h/d=hours per day; h/m=hours per month; h/w=hours per week; LogRA=logistic regression analysis; ns=no association; PSQ-I=Pittsburgh Sleep Quality Index; SDS=Self-Rating Depression Scale; Sig=significant; (+)=positive association; (-)=negative association]

Author	Participants	Survey	Outcomes (measures), Statistical analysis	Working hours	Results
Kobayashi et al, 2012 (35)	N=933, Japan, workers in manufacturing company, all male, age 19–70 years old, including shift workers (25.8%)	2009	Metabolic syndrome (Japanese criteria (2005), using blood samples, sphygmomanometer, and waist circumference was measured by medical staff), LogRA	Working ≥ 7 and $\leq 8^a$, >8 and ≤ 9 , >9 and ≤ 10 , >10 h/d	All participants: Sig (+) 2.32-fold (working >10 h/d) Participants without shift workers: ns Age stratified analysis ≥ 40 years or <40 years: ≥ 40 group: Sig (+): 2.02-fold (working >8 ≤ 9 h/d), 3.14-fold (working >10 h/d) ≥ 45 years or <45 years: ≥ 45 group: Sig (+): 2.82-fold (working >8 ≤ 9 h/d), 5.13-fold (working >10 h/d) ≥ 50 years or <50 years: ≥ 50 group: Sig (+): 3.33-fold (working >8 ≤ 9 h/d), 4.86-fold (working >10 h/d)
Nakashima et al, 2011 (36)	N=1510, Japan, white-collar workers in light metal products manufacturing company, all male, mean age 42.6 years old (range 18–59)	2004	Sleep (PSQ-I; PSQ-I global score, subscales of poor sleep quality, short sleep duration (≤ 6 hours/day), daytime dysfunction), LogRA	Number of over-time <26 h/m ^a , ≥ 26 and <40 , ≥ 40 and <50 , ≥ 50 and <63 , ≥ 63 h/m (working hours of contract: 40 h/w)	PSQ-I global score: Sig (+) 1.67-fold (overtime ≥ 50 <63 h/m) 1.87-fold (overtime ≥ 63 h/m) Poor sleep quality: ns Short sleep duration: Sig (+) 1.43-fold (overtime ≥ 26 <40 h/m) 1.51-fold (overtime ≥ 40 <50 h/m) 1.75-fold (overtime ≥ 50 <63 h/m) 3.68-fold (overtime ≥ 63 h/m) Daytime dysfunction: Sig (+) 1.82-fold (overtime ≥ 26 <40 h/m) 2.06-fold (overtime ≥ 40 <50 h/m) 2.36-fold (overtime ≥ 50 <63 h/m) 2.58-fold (overtime ≥ 63 h/m)
Nash et al, 2010 (37)	N=2999, Australia, doctors, male 71%	2007	Psychiatric morbidity (GHQ-28), hazardous alcohol use (AUDIT), LogRA	Working $<40^a$, 40–49, 50–59, ≥ 60 h/w	Psychiatric morbidity: Sig (+) 1.41-fold (working 50–59 h/w) 1.65-fold (working ≥ 60 h/w) Hazardous alcohol use: Sig (-) 0.67-fold (working ≥ 60 h/w)
Driesen et al, 2010 (38)	N=7217, Netherlands, employees representing almost all sectors and trades of the Dutch labour force from 45 companies and organizations, male 73%, mean age 41.9 years old	1998	Depressive mood (single question “Did you feel down every day over the last two weeks?” derived from Balansmeter), LogRA	Working 36–40 ^a , >40 h/w	Men: Sig (-) 0.74-fold (working >40 h/w) Women: ns
Virtanen et al, 2009 (28)	N=2436–2459, UK, civil servants, male 76%, mean age 52.2 years old	1997–1999 (phase 5)	Sleep (short sleeping hours (<7 hours/day), difficulty in falling asleep, frequent waking during the night, early waking, waking without feeling refreshed) (self-reported, Jenkins scale), LogRA	Working 35–40 ^a , 41–55, >55 h/w	Short sleeping hours: Sig (+) 1.39-fold (working 41–55 h/w) 2.25-fold (working >55 h/w) (linear trend ^b Sig (+): 1.39-fold) Difficulty in falling asleep: ns (linear trend ^b ns) Frequent waking: ns (linear trend ^b ns) Early waking: ns (linear trend ^b ns) Waking unrefreshed: ns (linear trend ^b ns)
Nagashima et al, 2007 (39)	N=715, Japan, chemical factory workers, all male, mean age 44.1 years old	2003	Depressive state (SDS), anxiety (CFSI), LogRA	Working $\leq 199^a$, 200–219, 220–239, 240–259, 260–279, ≥ 280 h/m	Depressive state: Sig (+) 2.75-fold (working 260–279 h/m) Anxiety: Sig (+) 2.28-fold (working 260–279 h/m) 2.51-fold (working ≥ 280 h/m)
Sekine et al, 2006 (40)	N=3556, Japan, civil servants, male 67%, mean age 42.7 (SD=10.2, range 20–65) years old, including shift workers (21%)	2003	Sleep quality (PSQ-I global score), LogRA	Working 7–9 ^a , 9–11, >11 h/d	All participants: Sig (+) 1.71-fold (working >11 h/d) Men only: Sig (+) 1.49-fold (working >11 h/d) Women only: Sig (+) 2.02-fold (working >11 h/d)

^a Reference working group

^b Linear trend = increasing 10-hour working time (per 10-hour increase)

Study F (29) evaluated the association between long working hours and cognitive function. The study design had characteristics of both a prospective cohort and cross-sectional study. Participants were followed for a mean period of 5.5 (range 3.9–7.1) years. Five cognitive tests were conducted and high scores represented good cognitive function. At baseline, only Mill Hill vocabulary test scores of participants in the groups working 41–55 hours/week and >55 hours/week were significantly lower than those of the reference group. On the other hand, phonemic fluency test scores showed a significantly positive linear trend with working hours. At follow-up, Alice Heim 4-I test (reasoning test) scores of participants in the group working >55 hours/week and vocabulary test scores of participants in the groups working 41–55 hours/week and >55 hours/week were significantly lower compared to the reference group. Changes in reasoning test scores between baseline and follow-up showed a significant decrease among participants in the groups working 41–55 hours/week and >55 hours/week compared to the reference group.

Tarumi et al (30) surveyed the association between long working hours and mental disorders [International Statistical Classification of Diseases and Related Health Problems, tenth revision (ICD-10): F00-99, G90] and circulatory disorders (ICD-10: I00-99). Information on working hours was collected every year for four years and averaged working hours were used in the analysis. Incidence was 6.6/10 000 and 15.7/10 000 person-months for mental and circulatory disorders, respectively. After adjusting for covariates, no statistical significance was found.

Nakanishi et al conducted a cohort study, and reported two separate outcomes in articles published in September (31) and May (32) 2001. In this review, for convenience, we treated these as two separate studies.

The September article (31) evaluated the association between long working hours and impaired fasting glucose (IFG), on the one hand, and type 2 diabetes mellitus (DM), on the other hand. IFG and type 2 DM were diagnosed according to the criteria of the American Diabetes Association (1997) using blood samples or self-reported hypoglycemic medication use. Incidence was 24.1/1000 person-years for IFG or type 2 DM (15.0/1000 person-years for IFG, 9.1/1000 person-years for type 2 DM). The results showed a significantly decreased risk of IFG or type 2 DM for participants in the group working ≥ 11.0 hours/day [adjusted relative risk (RR_{adj}) 0.50 (95% CI 0.25–0.98)]. No statistical significance was found for IFG, but a significantly lower risk was found for type 2 DM among participants working ≥ 11.0 hours/day [RR_{adj} 0.30 (95% CI 0.09–0.94)].

Nakanishi et al's other article (32) investigated the association between long working hours, hypertension

(HT), and borderline HT. HT was defined according to the WHO criteria (1978) as: systolic BP ≥ 160 mm Hg and/or diastolic BP ≥ 95 mm Hg or taking antihypertensive medication. Borderline HT was defined as a BP between HT and normotension (systolic BP < 140 mm Hg and diastolic BP < 90 mm Hg). The incidence of HT and borderline HT was 19.4/1000 and 85.3/1000 person-years, respectively. The results indicated a significantly decreased risk among participants working long hours. For HT, participants in the group working ≥ 11.0 hours/day had a significantly decreased risk [RR_{adj} 0.33 (95% CI 0.11–0.95)]. For borderline HT, participants in the groups working 10.0–10.9 hours/day and ≥ 11.0 hours/day had a significantly decreased risk [RR_{adj} 0.63 (95% CI 0.43–0.91) and RR_{adj} 0.48 (95% CI 0.31–0.74), respectively].

Kawakami et al (33) investigated the association between long working hours and non-insulin dependent DM (NIDDM, type 2 DM). NIDDM was diagnosed according to the WHO criteria (1980). An annual screening test using a urine sample was conducted for all participants. If glucosuria was found, fasting plasma glucose (FPG) was measured. Participants with high FPG (≥ 110 mg/dl) underwent a 75 g oral glucose tolerance test. The incidence for NIDDM was 1.95 /1000 person-years. The results indicated a significantly increased risk among participants working >50 hours/month of overtime [HR_{adj} 3.73 (95% CI 1.41–9.90)].

Shields (34) conducted a survey to investigate the association between long working hours and MDE, weight gain, smoking, alcohol consumption, and physical activity. The incidence of MDE was 3% and 5% for men and women, respectively. The Shields' study defined standard working hours as working ≥ 35 – < 41 hours/week, and long working hours as ≥ 41 hours/week. Results of MDE showed that only women working long hours had a significantly increased risk [OR_{adj} 2.2 (95% CI 1.1–4.4)]. For the other four outcomes, this study took changes in working hours both at baseline and follow-up into consideration. In the statistical analysis, the reference group was defined as those working standard hours both at baseline and follow-up. The incidence for unhealthy weight gain, increased daily smoking, increased alcohol consumption, and decreased physical activity was 10% for both men and women, 9% for men and 7% for women, 34% for men and 25% for women, and 43% for men and 41% for women, respectively. This study showed the results of several working patterns, however, in this review, we dealt with those results with a “long-long” working pattern (working long hours both at baseline and follow-up) similar to other cohort studies. A “long-long” working pattern was not significantly associated with these four outcomes compared to the reference group.

Cross-sectional design

Kobayashi et al (35) investigated the association between long working hours and metabolic syndrome using Japanese criteria. Overall, prevalence was 11.8% among participants. This study conducted statistical analyses with or without shift workers (25.8%). The model without shift workers indicated no significant association with long working hours. On the other hand, the adjusted model with a shift work schedule showed a significantly increased risk among participants working >10 hours/day [OR_{adj} 2.32 (95% CI 1.04–5.16)]. Several age-stratified analyses were conducted and indicated that those in the older age group with long worktime were at increased risk for metabolic syndrome compared to those of the same age in the reference group.

Nakashima et al (36) investigated the association between long working hours and sleep condition using the Pittsburgh Sleep Quality Index (PSQ-I). Working hours were recorded by timecards, and mean overtime of hours/month was calculated from data six months previously. A total of 35.8% of workers were above the PSQ-I cut-off point (poor sleepers). Poor sleepers and subscales of PSQ-I (poor sleep quality, short sleep duration, and daytime dysfunction) were analyzed. For poor sleepers, participants who worked ≥ 50 hours/month overtime had a significantly increased risk. The OR_{adj} were 1.67 (95% CI 1.17–2.38) for ≥ 50 and <63 hours/month overtime and 1.87 (95% CI 1.30–2.68) for working ≥ 63 hours/month overtime. Short sleep duration and daytime dysfunction were significantly associated with long working hours. The more overtime participants worked, the greater the OR_{adj} . Poor sleep quality was not significantly associated with the number of overtime hours worked.

Nash et al (37) surveyed the relationship between long working hours and psychiatric morbidity and hazardous alcohol use. Outcomes were measured by GHQ-28 and the Alcohol Use Disorders Identification Test (AUDIT). The prevalence of psychiatric morbidity was 28% and that of hazardous alcohol use was 15%. Participants working ≥ 50 hours/week were significantly more likely to suffer from psychiatric morbidity. The OR_{adj} were 1.41 (95% CI 1.05–1.89) for working 50–59 hours/week and 1.65 (95% CI 1.20–2.26) for working ≥ 60 hours/week. For hazardous alcohol use, only the group working ≥ 60 hours/week had a significantly decreased risk [OR_{adj} 0.67 (95% CI 0.45–0.99)].

Driesen et al (38) reported a survey from the Maastricht Cohort Study in 1998. In this review, we describe the association between long working hours and depressive mood according to our inclusion criteria. Depressive mood was assessed by the question, “Did you feel down every day over the last 2 weeks?” This wording was derived from the Balansmeter screening questionnaire. The prevalence was 6.8% for participants work-

ing 36–40 hours/week and 5.6% for those working >40 hours/week. The results were analyzed separately by gender because of a significant interaction between working hours and gender. A significant association was only found among men working >40 hours/week [OR_{adj} 0.74 (95% CI 0.56–0.99)].

Virtanen et al (28) investigated the association between long working hours and sleep condition. This study used baseline data from the cohort study described previously (studies D and E), and outcomes were the same. Only short sleeping hours were significantly associated with long working hours. The OR_{adj} were 1.39 (95% CI 1.16–1.68) for working 41–55 hours/week and 2.25 (95% CI 1.62–3.12) for working >55 hours/week.

Nagashima et al (39) reported a survey conducted on workers in 2003. In this review, we describe the association between long working hours, depressive state, and anxiety according to our inclusion criteria. Depressive state was measured by the Self-Rating Depression Scale (SDS) and anxiety was measured by the Cumulative Fatigue Symptom Index (CFSI). For depressive state, only participants in the group working 260–279 hours/month had a significantly increased risk [OR_{adj} 2.75 (95% CI 1.39–5.46)]. For anxiety, participants working ≥ 260 hours/month had a significantly increased risk, OR_{adj} 2.28 (95% CI 1.20–4.35) for working 260–279 hours/month and 2.51 (95% CI 1.29–4.90) for working ≥ 280 hours/month.

Sekine et al (40) reported the association between long working hours and sleep condition using PSQ-I. The prevalence of poor sleepers was 21.2% among men and 31.4% among women. All participants working >11 hours/day had a significantly increased risk [OR_{adj} 1.71 (95% CI 1.28–2.29)]. Gender stratified analyses showed a significant risk among both genders: men working >11 hours/day [OR_{adj} 1.49 (95% CI 1.03–2.15)] and women working >11 hours/day [OR_{adj} 2.02 (95% CI 1.24–3.31)].

Discussion

In this review, we explored articles eligible for inclusion published between 1995–2012. We found a total of 17 articles and 19 studies and described the association between long working hours and health. We identified all types of associations (positive, negative, or none).

Outcomes

The articles included in our review dealt with various outcomes, thus, the relationship of each with long working hours was investigated.

All-cause mortality. Only one prospective cohort study

(27) dealt with all-cause mortality. The results showed no significant association with long working hours. We could not reach any conclusion about this outcome due to the limited data available.

Circulatory disease. Four prospective cohort studies (24, 27, 30, 32) had six outcomes related to circulatory diseases. However, the results were inconsistent. Tarumi et al (30) indicated circulatory disorders including HT (ICD-10: I00-99) had no significant association with long working hours. We suspect that this result was affected by the small number of covariates (Appendix A: http://www.sjweh.fi/data_repository.php). Nakanishi et al (32) revealed that participants working long hours had a significantly decreased risk of HT and borderline HT. They described two reasons for these results: (i) whether the participants perceived overwork as stressful or not, and (ii) significantly higher 24-hour energy expenditure in the group working long hours. On the other hand, Nakamura et al (24) showed significantly increased diastolic BP in the group working ≥ 80.0 hours/month of overtime. We could not reach any conclusion regarding long working hours and BP.

The number of results related to CHD in our review were few and inconsistent (27, 30). However, in one meta-analysis about long working hours and CHD (41), these two articles (27, 30) were the basis for a subgroup analysis focusing on daytime workers. The results indicated an increased RR of 1.51 (95% CI 1.12–2.03) compared to those without long-term work. Thus, with the result of this meta-analysis, we concluded that working long hours is associated with CHD.

Diabetes mellitus. We found two studies (31, 33) that presented opposing results. Nakanishi et al (31) showed a significantly decreased risk associated with long working hours and the development of type 2 DM. However, Kawakami et al (33) indicated a significantly increased risk. Nakanishi et al (31) discussed the reasons for the discrepancy in great detail. They considered that it was caused by the difference in work environment and characteristics of participants, including the proportion of shift workers [0% (31) versus 46.3% (33)], the method of diagnosing DM, and high energy expenditure observed among participants working long hours (31). We agree with these suggestions, especially concerning the method of diagnosing of DM. Nakanishi et al (31) used blood samples and self-reported hypoglycemic medication use to diagnose DM, while Kawakami et al (33) used urine samples, blood samples, and 75 g OGTT. The latter would have been more accurate compared to the former. We also suspect that including shift workers among the participants made it more difficult to interpret the results.

We found one meta-analysis which investigated long working hours and type 2 DM (42), but the results showed no association. Therefore, we could not reach any conclusion about the association between long working hours and DM.

Metabolic syndrome. We found one cross-sectional study (35) about metabolic syndrome. The results of the statistical analysis changed depending on whether shift workers were included or not. Only the results with shift workers showed a significantly increased risk. As mentioned in the introduction section, shift work is detrimental to health. We believe that further studies are needed with regards to this outcome.

Depressive state. We found five studies (25, 26, 34, 38, 39) about depressive state. One cross-sectional study (38) indicated that long working hours significantly decreased the risk of depressive mood. We believe that this finding resulted from the way in which the outcome was measured. In the study, depressive mood was evaluated by a simple question, similar to one of the criteria for MDE in the DSM-IV-TR (Diagnostic and Statistical Manual of Mental Disorders, fourth edition, text revision) published by the American Psychiatric Association. However, it might be insufficient for measuring depressive mood. The reason is that patients themselves may not perceive they are experiencing a psychologically ill state, especially a depressive state. The other three prospective cohort studies (25, 26, 34) and one cross-sectional study (39) showed a significantly increased risk with long working hours. Consequently, we concluded that long working hours is associated with depressive state.

Anxiety. One prospective cohort study (26) and a cross-sectional study (39) investigated anxiety. Both results showed a significant increased risk for anxiety. We concluded that long working hours are indeed associated with anxiety. The results indicated that the risk rose among participants working >55 hours/week (26), 260–279 hours/month, or ≥ 280 hours/month (39). We suspect the existence of a threshold for developing anxiety.

Other mental states. We surveyed mental disorders (ICD-10: F00-99) in a prospective cohort study (30) and psychiatric morbidity in a cross-sectional study (37). We could not come to any conclusion about the association with long working hours due to the lack of data.

Sleep. We found five [two prospective cohort (28) and three cross-sectional (28, 36, 40)] studies that investigated sleep condition. A significant negative effect of long working hours on sleep condition was observed in all studies. We concluded that long working hours is associated with sleep condition.

In the two cohort studies (studies D and E) (28), study E identified more significant associations compared to study D. This might be attributable to the difference in measurement of working hours. Hence we consider that measuring working hours over a longer period is more effective to investigate the association between long working hours and health.

Two studies (36, 40) using PSQ-I to evaluate sleep condition presented similar results. This fact strongly supports the relationship between long working hours and sleep condition. All five studies had detailed subscales of sleep condition, and short sleeping hours or duration were highlighted regardless of the study design. Thus, among factors associated with sleep condition, short sleeping hours has been strongly associated with long working hours.

Cognitive function. We found one prospective cohort study (29) that evaluated cognitive function using five cognitive tests. However, one article was insufficient to enable us to reach any conclusion.

Behavior. We found one prospective cohort (34) and a cross-sectional (37) study examining behavior. Outcomes were alcohol use, physical activity, smoking, and weight gain. In the cohort study (34), no significant association was found between long working hours and the four outcomes. The cross-sectional study (37) showed a significantly decreased risk of hazardous drinking in the group working ≥ 60 hours/week. We inferred that the reason for this was the lack of personal time, which would in turn reduce the opportunity for drinking. From these results, we could not conclude any association between long working hours and health-related behavior.

However, one cohort study (34) revealed an important implication. In the results section, we only described the results of a long-long working pattern (working long hours both at baseline and follow-up) similar to other cohort studies. But participants with “standard-long” working patterns (working standard hours at baseline and long hours at follow-up) were at a significantly increased risk for alcohol consumption, smoking, and weight gain, compared to a “standard-standard” working pattern (working standard hours both at baseline and follow-up) (Appendix A, http://www.sjweh.fi/data_repository.php). Thus, a change in working hours from standard to long working hours could lead to an unhealthier lifestyle.

Summary and interpretation

We conclude that long working hours is associated with depressive state, anxiety, sleep condition, and CHD. Although compared to a previous review (23) we concluded that only a small number of health outcomes are associated with long working hours, we could clearly

base our results on epidemiological evidence due to our strict inclusion criteria. To be more precise, due to our strict inclusion criteria, we were able to exclude differences in the definition of long working hours and the detrimental effect of shift work. By using the standard working group as the reference group, we believe we could find clearer evidence for the association between long working hours and health.

We aimed to exclude differences in the definition of long working hours and the influence of shift work for several reasons. Concerning the former, we found disparities in the definition of long working hours in previous studies and recognized this to be problematic because the results of each study were not easily comparable. Consequently, this could confuse the interpretation of the association between long working hours and health. By excluding differences in the definition of long working hours, it is easier to compare previous studies with each other. Furthermore, by comparing and unifying the results, we could reach a universal conclusion about these associations. With regards to the influence of shift work, we considered that this factor itself raised a health risk for workers (17–19). This has been implied by one meta-analysis (41) that investigated long working hours and CHD. The overall RR was 1.80 (95% CI 1.42–2.29) for long working hours. As described above, a subgroup analysis that was limited to daytime workers and did not include shift workers showed an RR of 1.51 (95% CI 1.12–2.03), lower than the overall RR. The exclusion of the influence of shift work seemed to decrease the health risk for long working hours and was, therefore, important in the studies about the association between long working hours and health.

We suspected that the health risk for long working hours in previous studies might confuse our interpretation unless appropriate consideration of the definition of long working hours and the influence of shift work was carried out. We strongly suggest future research that takes this into account when clarifying the association between long working hours and health.

Reasons behind the exclusion criteria

We identified 149 articles after our initial search. However, after a fulltext review, only 17 articles remained. The most common reason for exclusion was “participant criteria” (N=70), which concerned shift work and the reference group. Though we understand that each study has its own purpose and survey items differ according to the purpose, due to its detrimental effect on health, we believe that shift work has to be considered in the study design when evaluating the association between long working hours and health. From our search, only Kobayashi et al (35) presented results with and without shift workers. The significant negative impact on health observed

among participants including shift workers disappeared in the results where shift workers were excluded. Even if statistical analyses are adjusted for work schedules, the influence of shift work may still partially remain.

Causal pathway

We have briefly inferred a causal pathway, which might explain why long working hours appear to increase the risk to health. Workers with long working hours need more time to recover from work (43). However, long working hours decrease the amount of private time available. Therefore, workers with long working hours do not have enough time to recover from exhaustion. The small amount of private time leads to an irregular lifestyle.

Such an irregular lifestyle may result in sleep problems and influence health-related behavior. If sleep problems, such as short sleeping hours and difficulty in falling asleep, continue for a given period, workers may develop lack of sleep, and insufficient sleep may lead to depression (44). Depression is a well-known risk factor for the development of CHD (45, 46). One meta-analysis (46) showed that subjects suffering from depression were at increased risk for developing CHD compared to non-depressed subjects. Short sleep time and frequent lack of sleep are also associated with an increased risk of acute myocardial infarction (7).

Irregular lifestyle caused by long working hours might also influence health-related behaviors such as alcohol consumption, smoking, and weight gain. From our search, we found that detrimental changes in health-related behaviors might occur when changes in working hours occur, such as changing from standard to long working hours (34). We suggest that these changes in health-related behaviors may cause a pre-morbid state, such as chronic liver or lung damage and the development of obesity, which in turn may lead to the development of diseases such as CHD and DM.

Limitations

The interpretation of our findings requires the consideration of several limitations. The first is selection bias. In our initial search, we used two databases with eight keywords. We selected peer-reviewed articles published in English with abstracts. Thus, articles that did not meet these requirements were not included in our review. The second limitation is publication bias. Although we examined many articles, there might be other studies that were not published because their results showed no significance. This might cause an overestimation about the relationship between long working hours and health. The third limitation is the measurement of working hours. Only two studies used timecards. We agree with the measurement of working hours using

questionnaires or interviews because, in some cases (31, 32), participants who hold managerial positions may not record their working hours using a time clock. However, self-reported working hours may also be inaccurate.

Concluding remarks

Our systematic review concluded that long working hours is associated with depressive state, anxiety, sleep condition, and CHD. Though the number of health outcomes we ascertained was small due to our strict inclusion criteria, we were able to exclude differences in the definition of long working hours and the influence of shift work. Since both factors may influence the association between long working hours and health, further studies are necessary that deal with them appropriately.

Acknowledgments

The authors greatly appreciate all persons who engaged in the related articles and replied to our inquiries. Both authors declare no conflict of interest.

References

1. Japanese Ministry of Health, Labour and Welfare [internet]. Tokyo: [updated 2002 May 22; cited 2012 Nov 7]. Available from: <http://www.mhlw.go.jp/houdou/2002/05/h0522-2.html>.
2. Japanese Ministry of Health, Labour and Welfare [internet]. Tokyo: [updated 2012 June 15; cited 2012 Nov 7]. Available from: <http://www.mhlw.go.jp/stf/houdou/2r9852000002coxc.html>.
3. OECD (Organisation for Economic Co-operation and Development). StatExtracts average annual hours actually worked per worker [internet]. Paris: OECD [cited 2012 Nov 7]. Available from: http://stats.oecd.org/Index.aspx?DatasetCode=AVE_HRS.
4. Cheng Y, Park J, Kim Y, Kawakami N. The recognition of occupational diseases attributed to heavy workloads: experiences in Japan, Korea, and Taiwan. *Int Arch Occup Environ Health*. 2012 Oct;85(7):791–9. <http://dx.doi.org/10.1007/s00420-011-0722-8>.
5. The International Labour Organization, Working time around the world [internet]. Geneva. c1996-2013 [cited 2013 March 31]. Available from: http://www.ilo.org/wcmsp5/groups/public/---dgreports/---dcomm/---publ/documents/publication/wcms_104895.pdf.
6. Holtermann A, Mortensen OS, Burr H, Sogaard K, Gyntelberg F, Suadicani P. Long work hours and physical fitness: 30-year risk of ischaemic heart disease and all-cause mortality among middle-aged Caucasian men. *Heart*. 2010 Oct;96(20):1638–44. <http://dx.doi.org/10.1136/hrt.2010.197145>.

7. Liu Y, Tanaka H; Fukuoka Heart Study Group. Overtime work, insufficient sleep, and risk of non-fatal acute myocardial infarction in Japanese men. *Occup Environ Med.* 2002 Jul;59(7):447–51. <http://dx.doi.org/10.1136/oem.59.7.447>.
8. Sasaki T, Iwasaki K, Oka T, Hisanaga N. Association of working hours with biological indices related to the cardiovascular system among engineers in a machinery manufacturing company. *Ind Health.* 1999 Oct;37(4):457–63. <http://dx.doi.org/10.2486/indhealth.37.457>.
9. Sasaki T, Iwasaki K, Oka T, Hisanaga N, Ueda T, Takada Y, Fujiki Y. Effect of working hours on cardiovascular-autonomic nervous functions in engineers in an electronics manufacturing company. *Ind Health.* 1999 Jan;37(1):55–61. <http://dx.doi.org/10.2486/indhealth.37.55>.
10. Iwasaki K, Sasaki T, Oka T, Hisanaga N. Effect of working hours on biological functions related to cardiovascular system among salesmen in a machinery manufacturing company. *Ind Health.* 1998 Oct;36(4):361–7. <http://dx.doi.org/10.2486/indhealth.36.361>.
11. Tomioka K, Morita N, Saeki K, Okamoto N, Kurumatani N. Working hours, occupational stress and depression among physicians. *Occup Med (Lond).* 2011 May; 61(3):163–70. <http://dx.doi.org/10.1093/occmed/kqr004>.
12. Schluter PJ, Turner C, Benefer C. Long working hours and alcohol risk among Australian and New Zealand nurses and midwives: a cross-sectional study. *Int J Nurs Stud.* 2012 Jun;49(6):701–9. <http://dx.doi.org/10.1016/j.ijnurstu.2012.01.005>.
13. Grosch JW, Caruso CC, Rosa RR, Sauter SL. Long hours of work in the U.S.: associations with demographic and organizational characteristics, psychosocial working conditions, and health. *Am J Ind Med.* 2006 Nov;49(11):943–52. <http://dx.doi.org/10.1002/ajim.20388>.
14. The Japan Institute for Labour Policy and Training [internet]. Tokyo: c2003-2013 [updated 2012 April 2; cited 2013 Mar 30]. Available from: http://www.jil.go.jp/kokunai/statistics/databook/2012/06/p195-202_16-6.pdf.
15. Nielsen NR, Stahlberg C, Strandberg-Larsen K, Kristensen TS, Zhang ZF, Hundrup YA, et al. Are work-related stressors associated with diagnosis of more advanced stages of incident breast cancers? *Cancer Causes Control.* 2008 Apr;19(3):297–303. <http://dx.doi.org/10.1007/s10552-007-9092-7>.
16. The International labour organization [Internet]. Geneva: c1996-2012 [cited 2012 Oct 10]. Available from: http://www.ilo.org/dyn/normlex/en/f?p=1000:12100:0::NO::P12100_INSTRUMENT_ID:312192.
17. Vyas MV, Garg AX, Iansavichus AV, Costella J, Donner A, Laugsand LE, et al. Shift work and vascular events: systematic review and meta-analysis. *BMJ.* 2012 Jul 26; 345:e4800. <http://dx.doi.org/10.1136/bmj.e4800>.
18. Knutsson A, Bøggild H. Gastrointestinal disorders among shift workers. *Scand J Work Environ Health.* 2010 Mar;36(2):85–95. <http://dx.doi.org/10.5271/sjweh.2897>.
19. Esquirol Y, Bongard V, Mabile L, Jonnier B, Soulat JM, Perret B. Shift work and metabolic syndrome: respective impacts of job strain, physical activity, and dietary rhythms. *Chronobiol Int.* 2009 Apr;26(3):544–59. <http://dx.doi.org/10.1080/07420520902821176>.
20. Megdal SP, Kroenke CH, Laden F, Pukkala E, Schernhammer ES. Night work and breast cancer risk: a systematic review and meta-analysis. *Eur J Cancer.* 2005 Sep; 41(13):2023–32. <http://dx.doi.org/10.1016/j.ejca.2005.05.010>.
21. Antunes Lda C, Jornada MN, Ramalho L, Hidalgo MP. Correlation of shift work and waist circumference, body mass index, chronotype and depressive symptoms. *Arq Bras Endocrinol Metabol.* 2010 Oct;54(7):652–6. <http://dx.doi.org/10.1590/S0004-27302010000700010>.
22. Wright KP Jr, Bogan RK, Wyatt JK. Shift work and the assessment and management of shift work disorder (SWD). *Sleep Med Rev.* 2013 Feb;17(1):41–54. <http://dx.doi.org/10.1016/j.smrv.2012.02.002>.
23. van der Hulst M. Long workhours and health. *Scand J Work Environ Health.* 2003; 29(3):171–88. <http://dx.doi.org/10.5271/sjweh.720>.
24. Nakamura K, Sakurai M, Morikawa Y, Miura K, Ishizaki M, Kido T, et al. Overtime work and blood pressure in normotensive Japanese male workers. *Am J Hypertens.* 2012 Sep;25(9):979–85. <http://dx.doi.org/10.1038/ajh.2012.37>.
25. Virtanen M, Stansfeld SA, Fuhrer R, Ferrie JE, Kivimäki M. Overtime work as a predictor of major depressive episode: a 5-year follow-up of the Whitehall II study. *PLoS One.* 2012;7(1):e30719. <http://dx.doi.org/10.1371/journal.pone.0030719>.
26. Virtanen M, Ferrie JE, Singh-Manoux A, Shipley MJ, Stansfeld SA, Marmot MG, et al. Long working hours and symptoms of anxiety and depression: a 5-year follow-up of the Whitehall II study. *Psychol Med.* 2011;41(12):2485–94. <http://dx.doi.org/10.1017/S0033291711000171>.
27. Virtanen M, Ferrie JE, Singh-Manoux A, Shipley MJ, Vahtera J, Marmot MG, et al. Overtime work and incident coronary heart disease: the Whitehall II prospective cohort study. *Eur Heart J.* 2010 Jul;31(14):1737–44. <http://dx.doi.org/10.1093/eurheartj/ehq124>.
28. Virtanen M, Ferrie JE, Gimeno D, Vahtera J, Elovainio M, Singh-Manoux A, et al. Long working hours and sleep disturbances: the Whitehall II prospective cohort study. *Sleep.* 2009 Jun;32(6):737–45.
29. Virtanen M, Singh-Manoux A, Ferrie JE, Gimeno D, Marmot MG, Elovainio M, et al. Long working hours and cognitive function: the Whitehall II Study. *Am J Epidemiol.* 2009 Mar 1;169(5):596–605. <http://dx.doi.org/10.1093/aje/kwn382>.
30. Tarumi K, Hagihara A, Morimoto K. A prospective observation of onsets of health defects associated with working hours. *Ind Health.* 2003 Apr;41(2):101–8. <http://dx.doi.org/10.2486/indhealth.41.101>.
31. Nakanishi N, Nishina K, Yoshida H, Matsuo Y, Nagano K, Nakamura K, et al. Hours of work and the risk of developing impaired fasting glucose or type 2 diabetes mellitus in Japanese male office workers. *Occup Environ Med.* 2001 Sep;58(9):569–74. <http://dx.doi.org/10.1136/oem.58.9.569>.

32. Nakanishi N, Yoshida H, Nagano K, Kawashimo H, Nakamura K, Tatara K. Long working hours and risk for hypertension in Japanese male white collar workers. *J Epidemiol Community Health*. 2001 May;55(5):316–22. <http://dx.doi.org/10.1136/jech.55.5.316>.
33. Kawakami N, Araki S, Takatsuka N, Shimizu H, Ishibashi H. Overtime, psychosocial working conditions, and occurrence of non-insulin dependent diabetes mellitus in Japanese men. *J Epidemiol Community Health*. 1999 Jun;53(6):359–63. <http://dx.doi.org/10.1136/jech.53.6.359>.
34. Shields M. Long working hours and health. *Health Rep*. 1999 Autumn;11(2):33–48 (English); 37–55 (French).
35. Kobayashi T, Suzuki E, Takao S, Doi H. Long working hours and metabolic syndrome among Japanese men: a cross-sectional study. *BMC Public Health*. 2012 May 31;12:395. <http://dx.doi.org/10.1186/1471-2458-12-395>.
36. Nakashima M, Morikawa Y, Sakurai M, Nakamura K, Miura K, Ishizaki M, et al. Association between long working hours and sleep problems in white-collar workers. *J Sleep Res*. 2011 Mar;20(1 Pt 1):110–6. <http://dx.doi.org/10.1111/j.1365-2869.2010.00852.x>.
37. Nash LM, Daly MG, Kelly PJ, van Ekert EH, Walter G, Walton M, et al. Factors associated with psychiatric morbidity and hazardous alcohol use in Australian doctors. *Med J Aust*. 2010 Aug 2;193(3):161–6.
38. Driesen K, Jansen NW, Kant I, Mohren DC, van Amelsvoort LG. Depressed mood in the working population: associations with work schedules and working hours. *Chronobiol Int*. 2010 Jul;27(5):1062–79. <http://dx.doi.org/10.3109/07420528.2010.489877>.
39. Nagashima S, Suwazono Y, Okubo Y, Uetani M, Kobayashi E, Kido T, et al. Working hours and mental and physical fatigue in Japanese workers. *Occup Med (Lond)*. 2007 Sep;57(6):449–52. <http://dx.doi.org/10.1093/occmed/kqm047>.
40. Sekine M, Chandola T, Martikainen P, Marmot M, Kagamimori S. Work and family characteristics as determinants of socioeconomic and sex inequalities in sleep: The Japanese Civil Servants Study. *Sleep*. 2006 Feb;29(2):206–16.
41. Virtanen M, Heikkilä K, Jokela M, Ferrie JE, Batty GD, Vahtera J, et al. Long working hours and coronary heart disease: a systematic review and meta-analysis. *Am J Epidemiol*. 2012 Oct 1;176(7):586–96. <http://dx.doi.org/10.1093/aje/kws139>.
42. Cosgrove MP, Sargeant LA, Caleyachetty R, Griffin SJ. Work-related stress and Type 2 diabetes: systematic review and meta-analysis. *Occup Med (Lond)*. 2012 Apr;62(3):167–73. <http://dx.doi.org/10.1093/occmed/kqs002>.
43. Jansen N, Kant I, van Amelsvoort L, Nijhuis F, van den Brandt P. Need for recovery from work: evaluating short-term effects of working hours, patterns and schedules. *Ergonomics*. 2003 Jun 10;46(7):664–80. <http://dx.doi.org/10.1080/0014013031000085662>.
44. Nakata A. Work hours, sleep sufficiency, and prevalence of depression among full-time employees: a community-based cross-sectional study. *J Clin Psychiatry*. 2011 May;72(5):605–14. <http://dx.doi.org/10.4088/JCP.10m06397gry>.
45. Sundquist J, Li X, Johansson SE, Sundquist K. Depression as a predictor of hospitalization due to coronary heart disease. *Am J Prev Med*. 2005 Dec;29(5):428–33. <http://dx.doi.org/10.1016/j.amepre.2005.08.002>.
46. Rugulies R. Depression as a predictor for coronary heart disease. a review and meta-analysis. *Am J Prev Med*. 2002 Jul;23(1):51–61. [http://dx.doi.org/10.1016/S0749-3797\(02\)00439-7](http://dx.doi.org/10.1016/S0749-3797(02)00439-7).

Received for publication: 17 December 2012